

Combat-Related Polytrauma (non-live)

Goals and Objectives

Course Description

“Combat-Related Polytrauma” is an online recorded video of a previously presented live CE webinar for physical therapists and physical therapist assistants. This course presents contemporary information about somatic trauma and comorbidities in combat veterans. This course includes a review of current literature relating to the pathophysiology, clinical presentation, and therapeutic management.

Course Rationale

The purpose of this course is to provide participants with contemporary information about conflict-related multiple trauma. Physical therapy professionals can use this information to develop and implement effective treatment programs that address the specific needs of individuals effected by these disorders.

Course Goals and Objectives

Upon completion of this course, participants will be able to:

1. List and describe common combat-related traumatic injuries and conditions.
2. Detail comorbidities contributing to polytrauma.
3. Identify and define the roles of all health care professionals providing comprehensive care to individuals with combat-related polytrauma.
4. Detail the Department of Defense Standard Case Definitions for traumatic brain injury.
5. Identify the mechanism of injury, pathophysiology, and medical management of blast related traumatic brain injury.
6. Identify the mechanism of injury and rehabilitation considerations related to dismantled complex blast injury.
7. Describe common pain syndromes associated with combat-related polytrauma.
8. Identify rehabilitation considerations for non-somatic for disorders presenting post-deployment.
9. Identify and select appropriate screening tools to assess physical and psychosocial function.
10. Identify the clinical presentation and rehabilitative management of individuals presenting with polytrauma.

Course Provider – Innovative Educational Services

Course Instructor - Jodi Gootkin, PT, Med, CEAS

Target Audience – Physical Therapists, Physical Therapist Assistants

Course Educational Level – This course is applicable for introductory learners.

Course Prerequisites – None

Method of Instruction/Availability – Recorded video available online on-demand

Criteria for Issuance of CE Credits – Completion of viewing of 3 hour recorded video, and a score of 70% correct or greater on the course post-test

Continuing Education Credits – Three (3) hours of continuing education credit.

Course Fee - \$34.95

Conflict of Interest – No conflict of interest exists for the presenter or provider of this course.

Refund Policy - Unrestricted 100% refund upon request. The request for a refund by the learner shall be honored in full without penalty or other consideration of any kind. The request for a refund may be made by the learner at any time without limitations before, during, or after course participation.

COMBAT RELATED POLYTRAUMA

1

Live Interactive Webinar Presented By:
Jodi Gootkin, PT, MEd, CEAS
jodiemail@comcast.net

Copyright J. Gootkin 2019

Course Overview

2

“Combat Related Polytrauma” is a live (real-time) interactive webinar for rehabilitation professionals that presents contemporary information about somatic trauma and comorbidities in combat Veterans. This course includes a review of current literature relating to the pathophysiology, clinical presentation, and therapeutic management.

Copyright J. Gootkin 2019

Course Rationale

3

The purpose of this course is to provide participants with contemporary information about conflict-related multiple trauma. Rehabilitation professionals can use this information to develop and implement effective treatment programs that address the specific needs of individuals effected by these disorders.

Copyright J. Gootkin 2019

Goals and Objectives

4

1. Detail comorbidities contributing to polytrauma.
2. List and describe common battlefield exposures that result in combat-related trauma injuries.
3. Identify the mechanism of injury, pathophysiology, and medical management of blast related injury.
4. Detail considerations for successful therapeutic interactions with a veteran transitioning to civilian rehabilitation settings.
5. Identify medical management and research advances for combat related amputations and spinal cord injuries.
6. Detail the Department of Defense Standard Case Definitions for traumatic brain injury.
7. Describe the clinical presentation and interaction strategies for an individual with post traumatic stress disorder.
8. Identify common pain syndromes associated with combat-related polytrauma.
9. Identify rehabilitation considerations for non-somatic disorders presenting post-deployment.
10. Identify appropriate screening tools to assess physical and psychosocial function.

Copyright J. Gootkin 2019

Disclaimer

5

Application of concepts presented in this webinar is at the discretion of the individual participant in accordance with federal, state, and professional regulations.

Copyright J. Gootkin 2019

“And I will always do my duty
No matter what the price
I've counted up the cost
I know the sacrifice”

~Chuck Cannon and Toby Keith
American Soldier Lyric

Copyright J. Gootkin 2019

United States Military Conflicts

9



Copyright J. Gootkin 2019

https://www.va.gov/vetsinworkplace/docs/em_datesnames.asp

Impact of Military Conflicts

10

“The legacy of this conflict will not only be what we have learned, but also how rapidly we were able to disseminate, educate, and change practice on the battlefield in nearly “real-time”

Brian Eastridge, MD, FACS
Colonel, MC, US Army
Trauma Consultant, US Army Surgeon General
<http://www.cs.amedd.army.mil/borden/book/ccc/UCLAFrontMatter.pdf>

Copyright J. Gootkin 2019

Survivorship

11

□ Improved body armor, surgical care in proximity to the battlefield, and faster evacuation have contributed to a greater than 90% combat survival rate.

Copyright J. Gootkin 2019

<http://www.apta.org/TBI/WoundedWarriors/>
<http://specialoperationsmedicine.org/Pages/tccc.aspx>

The casualty survival rate in Iraq and Afghanistan is the best in US history.

12



Copyright J. Gootkin 2019

<http://armymedicine.mil/Documents/DCBI-Task-Force-Report-Redacted-Final.pdf>

Our Veterans

13

- Since 2002 1.97 million Veterans have become eligible for VA benefits.
 - 61% Former Active Duty
 - 39% Reserve and National Guard
- Most common out-patient diagnoses
 - Musculoskeletal ailments
 - Mental disorders
 - Ill defined conditions

Epidemiology Program, Post-Deployment Health Group, Office of Patient Care Services, Veterans Health Administration, Department of Veterans Affairs. (2017). Analysis of VA Health Care Utilization among Operation Enduring Freedom, Operation Iraqi Freedom, and Operation New Dawn Veterans, from 1st Qtr FY 2002 through 3rd Qtr FY 2015. Washington, DC: Author

Copyright J. Gootkin 2019

Combat Related Polytrauma Definition

14

- The US Veterans Administration (VA) defines Polytrauma as injury to two or more body systems resulting in physical, cognitive, psychological, or psychosocial impairments and functional disability.
- Management requires a high level of integration and coordination of clinical care and with additional support services.

<https://www.polytrauma.va.gov/news-and-resources/terminology-and-definitions.asp>

Consider This

Copyright J. Gootkin 2019

Combat Related Injuries and Conditions

15

- Injuries are sustained during active combat or duty from Explosive Weapons and Improvised Explosive Devices resulting in physical, cognitive, psychological or psychosocial impairments.
 - Traumatic Brain Injury (TBI)
 - Amputation
 - Spinal Cord Injury (SCI)
 - Visual and hearing impairment
 - Burns
 - Musculoskeletal injuries
 - Post-Traumatic Stress Disorder

VA, Polytrauma Rehabilitation Center Design Guide 2014
<https://www.ctm.va.gov/tbl/dGuide/dgPRC.pdf>

Consider This

Copyright J. Gootkin 2019

Exposure Risks

- A variety of environmental and chemical exposures carry potential health risks contributing to casualties in combat.

- Sand, Dust, Particulate Matter
- Burn Pits
- Infectious Diseases
- Vibration
- Noise
- Depleted Uranium
- Blast Injuries and Toxic Embedded Fragments

Copyright J. Gootkin 2019

Exposure: Burn Pits

17

- Toxins released from burning waste may affect the skin, eyes, respiratory, cardiovascular, gastrointestinal, and other body systems.
- Effects are primarily temporary with limited evidence of association between exposure and reduced pulmonary function.

Baird, C. P. (2012). Review of the Institute of Medicine report: long-term health consequences of exposure to burn pits in Iraq and Afghanistan. ARMY PUBLIC HEALTH COMMAND ABERDEEN PROVING GROUND MD.

Respiratory Condition

COPD

Emphysema

Chronic Bronchitis

Copyright J. Gootkin 2019

Exposure: Infectious Disease

18

Copyright J. Gootkin 2019
<https://www.publichealth.va.gov/exposures/infectious-diseases/index.asp>

Exposure: Vibration

19

Hand-Arm Vibration Syndrome
Develops from continuous use of pneumatic, electric, hydraulic or gasoline powered hand tools.

➔

Whole Body Vibration
Impacts the musculoskeletal system when operating heavy equipment, trucks, helicopters and ships.

Copyright J. Gootkin 2019

Hand Arm Vibration Syndrome (HAVS)

20

- ❑ Vibration trauma damages the endothelial layer of hand capillaries, nerves, and tendons.
- ❑ Symptoms include
 - ❑ Blanching of one or more fingers with cold exposure
 - ❑ Tingling, numbness, decreased sensation in fingers
 - ❑ Hand pain
 - ❑ Decrease grip strength and dexterity

Copyright J. Gootkin 2019

Whole Body Vibration

21

- ❑ Low frequency vibration is transmitted through the body while sitting, standing, or reclined on a surface results in amplification of the vibration.
- ❑ Voluntary and involuntary muscular contraction occurs in response inducing fatigue and musculoskeletal damage particularly in the spine.

Copyright J. Gootkin 2019

Exposure: Noise

22

- ❑ Gunfire, explosions, rockets, heavy weapons, aircrafts, and machinery create high intensity noise and vibrations leading to hearing loss and tinnitus.
- ❑ Sound waves are directed into the cochlea and can't stop crushing the hair cells that are responsible for high-pitched hearing such as voices.

Impulse Impact Noise	Less than ½ a second	Explosions, weapon fire, grenades	Sudden or gradual hearing loss
Steady State Noise	Longer than ½ a second	Tactical vehicles, generators, aircraft	Gradual loss worsening over time

Image: http://www.public.navy.mil/navsafecen/Pages/acquisition/vibration_acquisition.aspx

Copyright J. Gootkin 2019

Exposure: Depleted Uranium

23

- ❑ Manufacturing of enriched uranium for nuclear reactors or weapons yields depleted uranium (DU) as a byproduct.
- ❑ It possesses 60% the radioactivity and same chemical toxicity as natural uranium.
- ❑ DU is utilized in tank armor and some bullets for better penetration of armored vehicles.
- ❑ When this type of projectile penetrates the vehicle some of the pieces of DU can become embedded in soft tissue, inhaled or swallowed.

Image: https://www.publichealth.va.gov/exposures/toxic_fragments/index.asp

Copyright J. Gootkin 2019

Depleted Uranium

24

- ❑ VA currently reports no health complications in Veterans.
- ❑ WHO notes that further research is needed to determine the extent, reversibility, and thresholds for organ damage with exposure.

<p style="text-align: center; margin: 0;">Inhaled</p> <ul style="list-style-type: none"> • Lung irradiation damage • Lung cancer in high doses 	<p style="text-align: center; margin: 0;">Embedded Fragments</p> <ul style="list-style-type: none"> • Uranium detectable in urine without accompanying medical compromise • Possible kidney damage if high doses
---	---

Image: https://www.publichealth.va.gov/docs/depleted-uranium/du_factsheet.pdf
http://www.who.int/ionizing_radiation/pub_meet/en/DU_Enq.pdf

Copyright J. Gootkin 2019

Exposure: Blast Injury 25

□ Improvised Explosive Devices create violent explosions containing metal or composite foreign material projectiles spread over a large radius.

Plastic	Ceramic	Glass	Wood
Cement	Nails	Metal	Rocks

□ Heat, pressure waves, and fragments cause large zones of soft tissue injury and damage to multiple body systems.

Copyright J. Gootkin 2019

Classifications of Blast Injury 26

- Primary Injury Barotrauma Damage**
 - Initial shock wave damages hollow organs.
 - Lungs, ears, gastrointestinal tract, eyes, brain, traumatic amputation
- Secondary Injury Fragmentation Injuries**
 - Penetrating wounds from debris propelled by shock waves.
 - Any region of the body, soft tissues, eyes
- Tertiary Injury Impact Injuries**
 - Displacement of victim by blast force.
 - Fractures, amputation, head/neck trauma
- Quaternary Injury Indirect Trauma**
 - Thermal burns, inhalation of gas/debris, crush injuries, exacerbation of pre-existing conditions
- Quinary Injury Contamination**
 - Biological, chemical, radioactive contamination

Copyright J. Gootkin 2019

Dismounted Complex Blast Injury 27

□ During counter-insurgency efforts, increased use of foot patrols led to the emergence of multiple limb amputations, complex pelvic/perineal wounds, spin injuries, and multiple penetrating fragment wounds.

□ By 2010, massive transfusion requiring 16 to over 25 units of bloods, major limb amputation, genitourinary injuries and traumatic brain injury rates increased drastically.

Wallace, D. (2012). Trends in traumatic limb amputation in Allied Forces in Iraq and Afghanistan. *Journal of Military and Veterans Health*, 20(2), 31. <http://armymedicine.mil/Documents/DCBI-Task-Force-Report-Redacted-Final.pdf>

Copyright J. Gootkin 2019

Exposure: Toxic Embedded Fragments 28

□ Improvised Explosive Device and blast shrapnel, bullets, or other projectiles cause injury where they enter the body.

□ Secondary pathology results as the chemicals embedded in the fragment that has not been removed from the body dissolve traveling to distant regions through the bloodstream inducing pathology.

□ According to the VA “No one knows the specific health problems that might happen because of the fragments”

<https://www.publichealth.va.gov/docs/exposures/TEFSC-Veterans-fact-sheet.pdf>
Image <https://www.publichealth.va.gov/exposures/toxic-fragments/index.asp>

Copyright J. Gootkin 2019

Tactical Combat Casualty Care 29

□ Initial battle field treatment to manage the effects of the various exposure risks is focused on temporary measures and evacuation.

□ Suppression of enemy fire and moving casualties to cover are the major concerns.

Copyright J. Gootkin 2019

Tactical Trauma Care Setting 30

□ Care is initially provided by non-medical combatant or combat lifesaver in the field.

Copyright J. Gootkin 2019

Tourniquets Save Lives

31

Timeline of tourniquet use:

- WW II 1945:** Strap and buckle rarely controlled bleeding.
- Vietnam 1970:** 2500 deaths from preventable hemorrhage.
- Gulf War 2004:** Still using makeshift tourniquets.
- Today:** Tourniquet carried on battle gear.

□ Appropriate placement of a tourniquet in the field before the onset of shock has been identified as a primary factor to minimize preventable death from hemorrhage.

Copyright J. Gootkin 2019 <http://www.cs.amedd.army.mil/borden/book/ccc/UCLAFrontMatter.pdf>

Modern Tourniquet

32

- **Combat Application**
Tourniquet is now standard issue equipment.
- **Removal within 6 hours** saves lives without increased incidence of secondary amputation due to ischemia.

Copyright J. Gootkin 2019 <https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFull.pdf>

Echelons of Care

33

- The Echelon structure refers to the level of command and control that is available to provide increasing levels of medical care.
- The goal is for the patient to be evacuated from the battle field and reach surgical care within one hour of injury.

Copyright J. Gootkin 2019

Continuum of Care

34

Continuum of Care:

- Point of Injury
- Echelon I: Care Under Fire, Tactical Field Care, Tactical Evacuation Care
- Echelon II: Forward Surgical Team
- Echelon III: Combat Support Hospital
- Echelon IV: Facility Outside Combat Zone
- Echelon V: Stateside (CONUS) Facilities
- VA and Civilian Rehabilitation

Copyright J. Gootkin 2019

Level I Care Under Fire

35

In battlefield with limited resources

Copyright J. Gootkin 2019

Level II Forward Surgical Team

36

Care facility removed from direct conflict with lab, imaging technology, surgical team.

Copyright J. Gootkin 2019

**Level III
Combat Support Hospital** 37

Field hospital with highly trained providers, labs, blood bank, operating rooms, imaging capabilities

Copyright J. Gootkin 2019

**Level IV
Critical Care Aeromedical Transport Team (CCAT) to
Out of Combat Zone Medical Center** 38

Major medical centers with capacity to diagnose and treat complex injuries. Example: Landstuhl, Germany

Copyright J. Gootkin 2019

**Level V -
US Based Medical Facility** 39

Fixed military medical facility close to home base or delivering most appropriate specialty care center.

Copyright J. Gootkin 2019

**Stateside Medical Care
Veterans Administration (VA)
Service Networks** 40

- For soldiers not returning to active duty, services are provided through the VA System.
- The country is subdivided into regions where various hospitals and outpatient service centers are located.

Copyright J. Gootkin 2019

VA Hub and Spoke Model of Care 41

□ This model emphasizes rehabilitation and a comprehensive continuum of care with full interdisciplinary teams of experts to manage specific chronic diagnoses.

Copyright J. Gootkin 2019

Veterans Choice Program 42

- The Veterans Choice Program enacted in 2014 with extended funding in 2017 expands the availability of medical services for eligible Veterans with community providers as opposed to waiting for an appointment with or traveling to a VA facility.
 - VA cannot schedule appointment within 30 days
 - Veteran resides more than 40 miles from closest VA facility.

Consider becoming a provider!

Copyright J. Gootkin 2019

Benefits of Veterans Choice Participation

43

❑ Veterans Choice community health providers have access to the VA electronic health record (EHR) that integrates all elements of a patient's health history, including medications, lab work, x-rays, scans, EKGs, medical diagnoses, and more to allow comprehensive coordinated care.

Copyright J. Gootkin 2019

Healthcare Team

44

- ❑ Determine the other members of the healthcare team who are providing care for comorbidities and recognize opportunities for additional referrals.
- ❑ Case Manager is typically the primary point of contact to coordinate services resources, and communication among the patient, family, military and VA.
- ❑ Benefits Counselor identifies VA system resources to help with education, employment, health, and financial support.

Copyright J. Gootkin 2019

Transitioning to Civilian Rehabilitation Settings

45

❑ Providing therapy to Veterans is a balance of managing their expectations with working through rehab ensuring clinical interactions are tailored to this unique community of patients.

Copyright J. Gootkin 2019

Military Cultural Competency

46

❑ Community providers must understand basic military culture to understand the unique needs of Veterans and build a strong therapeutic alliance.

70% VA providers

24% Community providers

8% General practitioners

Tantelian, Terri, Coreen Farris, Caroline Batka, Carrie M. Farmer, Eric Robinson, Charles C. Engel, Michael Robbins and Lisa H. Jaycox. Ready to Serve: Community-Based Provider Capacity to Deliver Culturally Competent, Quality Mental Health Care to Veterans and Their Families. Santa Monica, CA: RAND Corporation, 2014.

Copyright J. Gootkin 2019

Military Ethos

47

❑ "The overarching values of military culture set aside personal needs in favor of: teamwork, selflessness, shared goals, and perhaps most central, being of service to others - measuring one's worth by how well one serves others selflessly."

https://www.mentalhealth.va.gov/communityproviders/military_culture.asp

Copyright J. Gootkin 2019

War Zone vs. Civilian Life Culture Influences

48

War Zone	Civilian Life
Physically unsafe	Physically safe
Act, then think	Think, then act
Unpredictable	Predictable with routines
Follow orders	Cooperation and compromise
Numb or control emotions	Express feelings
Avoid closeness	Create intimacy and trust

WW II → Vietnam → Gulf War → OEF/OIF

- Immensely supported
- Soldiers shamed and dishonored
- Greeted as heroes
- Honored

Copyright J. Gootkin 2019

Transitioning Considerations for the Clinician 49

- Military service members transitioning into work, home life, medical and social interactions are negotiating many relationships and new situations.
- Consider military identity during your interactions.

Strength	Courage
Commitment	Loyalty
Accountability	Discipline
Persistence	Punctuality
Respect	

Copyright J. Gootkin 2019

Managing Clinical Interactions 50

- When integrating into civilian life hypervigilance, irritability, difficulty concentrating, and stress from personal relationships impacts interactions.
- Consider emotional health when determining treatment setting, scheduling, intervention selection, and home exercise program compliance.
- Building a strong therapeutic alliance fosters a relationship that encourages openness to additional resources for support.

Copyright J. Gootkin 2019

Do

- Thank Veteran for their service
- Ask about branch of service
- Talk about life outside the military
- Let Veteran share their experiences when they feel ready

51

Don't

- Discuss politics
- Ask about deaths or killing or "how it was over there"
- Assume they experienced combat
- Act like you know about their military career and experiences

Copyright J. Gootkin 2019

Social Services Referrals 52

- During our time with patients we have the opportunity for them to share multiple aspects of their personal circumstances that may lead to referrals for other services in addition to medical care.
 - Home Situation
 - Employment
 - Mental Health

Copyright J. Gootkin 2019

Social Service: Mental Health 53

- Encouraging Veterans to tracking their own psychological status and response to daily events aids in understanding symptoms and directing treatment.
- Department of Defense National Center for Telehealth and Technology developed the T2 MoodTracker App for service members to track behavior changes after combat deployments.
- Maintain an open dialogue and refer to mental health providers and self-help resources.

Copyright J. Gootkin 2019 <http://t2health.dcoe.mil/apps/t2-mood-tracker>

Social Services: Homelessness 54

- US Housing and Urban Development (HUD) estimates that on any single night in January 2016 40,000 Veterans were homeless and 13,000 were unsheltered or on the street.
- As of August 28, 2017 3 states and 50 communities have ended Veteran homelessness.

Copyright J. Gootkin 2019 https://www.va.gov/homeless/Homeless_Programs_General_Fact_Sheet_JAN_2017.pdf
<https://www.va.gov/HOMELESS/endingVetshomelessness.asp>

Homelessness Resources 55

- VA provides free resources for community outreach and education.
- Consider hanging poster in your waiting room.

https://www.va.gov/HOMELSS/get_involved.asp

Copyright J. Gootkin 2019

Social Services: Return to Employment 56

- Our interactions may highlight challenges the Veteran is experiencing in securing employment to provide for their family financially in addition to overall well being with personal satisfaction.
- Specific organizations are dedicated to assisting Veterans transition into the civilian workforce by offering career services, computer skills training, transition support, and job placements.

Copyright J. Gootkin 2019 <http://www.projecthired.org/jobseekers/Veterans>

Comprehensive Screening Considerations in Rehabilitation 57

- In addition to traditional initial evaluation procedures include questions and assessments to provide additional context to current injuries, promote a stronger therapeutic relationship, increase awareness of additional health conditions that may contribute to rehabilitation outcomes.
 - Military Service History
 - Respiratory Disease
 - Acoustic Trauma
 - Visual Impairments

Consider This

Copyright J. Gootkin 2019

Military Service History Interview 58

- The American Medical Association urges clinicians to obtaining information regarding military service that will help understand the unique medical conditions and potential concerns of Veterans.
- VA Healthcare provides guiding questions on the Military Health History Pocket Card.
 - Would it be ok if I talked with you about your military experience?
 - Were you or wounded, injured, or hospitalized?
 - Would it be ok to talk about your living situation?
 - Would it be ok if I asked about some things you may have been exposed to during your service?

Consider This

Copyright J. Gootkin 2019 <https://www.va.gov/oa/pocketcard/Military-Health-History-Card-for-print.pdf>

Military Terms 59

Awareness of some basic terminology a Veteran may use to describe their military experience will enhance communication.

Term or Acronym	Meaning
MOS Military Occupational Specialty	Job or career (infantryman, intelligence analyst, military police, etc.)
OCONUS	Outside Continental United States
CONUS	Inside Continental United States
Boots on the ground	To physically be in a location
Down range	Physically in a combat zone
In-country	Physically in a war zone
Theater	Geographic area directly involved in major operations involving combat

Copyright J. Gootkin 2019

Medical Surveillance of Embedded Fragments 60

- Superficial fragments that may work their way out of the body should be monitored and if removed sent to the VA Toxic Embedded Fragments Registry for analysis.
- Patients with suspected toxicity from fragments and depleted uranium should be referred for urine biomonitoring to assess concentrations of metals.

Arsenic	Cadmium	Chromium	Cobalt	Copper
Iron	Lead	Manganese	Nickel	Tungsten
Uranium	Zinc	Molybdenum		

Copyright J. Gootkin 2019 <http://www.cs.amedd.army.mil/FileDownloadPublic.aspx?docid=0781f05c-f914-4620-9598-66e52e9af4d7>

Respiratory Disease Assessment 61

- ❑ Combined exposures to dust storms, burn pits, and toxic chemicals may increase post deployment respiratory illness.
 - ❑ Asthma
 - ❑ COPD
 - ❑ Interstitial Lung Disease
- ❑ Suspected Tuberculosis requires chest x-ray.
- ❑ Monitor patients for decreased endurance and functional capacity.
 - ❑ Two or Six Minute Walk Test

Pugh, M. J., Jaramillo, C. A., Leung, K. W., Faverio, P., Fleming, N., Mortensen, E., ... B. Morris, M. J. (2016). Increasing Prevalence of Chronic Lung Disease in Veterans of the Wars in Iraq and Afghanistan. *Military medicine*, 181(5), 476-485.
 Sharkey, J. M., Abraham, J. H., Clark, L. L., Rohrbeck, P., Ludwig, S. L., Hu, Z., & Baird, C. P. (2016). Postdeployment respiratory health care encounters following deployment to Kabul, Afghanistan: a retrospective cohort study. *Military medicine*, 181(3), 265-271.

Copyright J. Gootkin 2019

Asthma vs. Tuberculosis (TB) 62

Asthma	Tuberculosis
Bilateral	Unilateral
Wheezing	Rales on auscultation
Non-productive cough	Productive cough >3 weeks
Triggers	Malaise and Fever
Relief with treatment	Night sweats

Copyright J. Gootkin 2019

Acoustic Trauma 63

- ❑ Barotrauma can rupture the tympanic membrane with small perforations healing spontaneously.
- ❑ Noise exposure or blast injury can cause vestibular impairment, otalgia, balance deficits, and hearing loss.

Distance from Blast	Primary Blast Injury	Secondary Blast Injury
0 - 50 feet	Fatal, eardrum rupture	Fatal
50 - 80 feet	Eardrum rupture	Fatal
80 - 130 feet	Temporary hearing threshold shift	Injury
130 - 1800 feet	None	Injury

<https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFULL.pdf>

Copyright J. Gootkin 2019

Auditory Impairments 64

- ❑ Assess patient for symptoms of compromise to the auditory system with referral to audiologist or otolaryngologist for additional evaluation.
 - ❑ Loss of ability to hear speech when background noise is present.
 - ❑ Diminished localization of hearing to distinguish direction sounds are coming from.
- ❑ Tinnitus
- ❑ Vertigo

Copyright J. Gootkin 2019

Visual Impairments 65

- ❑ Visual impairments range from low vision to total blindness from direct combat injuries or age-related diseases including macular degeneration, diabetic retinopathy, and glaucoma.
- ❑ Refer to VA Blind Rehabilitation Services for training in visual motor and perceptual skills, instruction in optical and electronic devices, and sensory training classes to promote safety and independence.

https://www.prosthetics.va.gov/blindrehab/BRS_Coordinated_Care.asp

Copyright J. Gootkin 2019

Vision Assessment 66

- ❑ Common complaints include blurred vision, photosensitivity, and accommodative problems.

```

    graph TD
      SRP[Self Reported Problems] --> FNA[Far-Near Acuity]
      FNA --> R[Reading]
      EABV[Eye Alignment and Bi-ocular Vision] --> C[Convergence]
      C --> A[Accommodation]
      S[Saccades] --> P[Pursuits]
      P --> VF[Visual Fields]
  
```

Radomski, M. Y., Finkelstein, M., Llanos, I., Scheiman, M., & Wagner, S. G. (2014). Composition of a Vision Screen for Servicemembers With Traumatic Brain Injury: Consensus Using a Modified Nominal Group Technique. *The American Journal of Occupational Therapy*, 68(4), 405-409.

Copyright J. Gootkin 2019

Vision Assessment: Convergence Insufficiency Symptom Survey

67

- ❑ Patients may have difficulty describing symptoms or may not realize fatigue, poor attention, dizziness, headache, balance deficits are related to vision impairment.
- ❑ Self assessment of how eyes feel when reading and doing close work.
- ❑ Score greater than 21 can indicate impairment requiring referral to ophthalmologist or optometrist

Copyright J. Gootkin 2019 <http://www.sankaranethralaya.org/pdf/patient-care/Convergence-Insufficiency-Symptom-Survey.pdf>

Vision Assessment: Snellen Eye Chart

68

- ❑ Widely used test of visual acuity.
- ❑ Patient is 20 feet away from chart
- ❑ Cover one eye
- ❑ Score is best line that they can read.
- ❑ Numbers on left side of chart indicate distance someone with normal vision can see that line at.
- ❑ Vision 20/40 or worse warrants referral to optometrist.

Copyright J. Gootkin 2019

Vision Assessment: Chronister Pocket Acuity Chart

69

- ❑ Small packet of numbers that are viewed from close distances too assess visual acuity from 20/200 to 20/10.
- ❑ Contains conversion chart to convert to Snellen values from multiple distances.
- ❑ Pediatric version available with image targets.
- ❑ Vision 20/40 or worse warrants referral to optometrist.

Copyright J. Gootkin 2019

Vision Assessment: Amsler Grid

70

- ❑ Tests metamorphopsia to detect alterations to central visual field requiring referral to ophthalmologist.
- ❑ Determines regions of visual field deficit by marking areas of wavy, distorted, broken or missing lines.

Copyright J. Gootkin 2019

Low Vision Patient Education

71

- ❑ Promote safety by recommending effective lighting, removal of clutter, installation of grab bars to avoid hazards.
- ❑ Modify the activities with education, training, and adaptive equipment.
 - ❑ Use contrast to improve visibility
 - ❑ Use overhead lighting to decrease shadows
 - ❑ Sit with back toward window or lamp

Copyright J. Gootkin 2019 <https://www.aota.org/-/media/Corporate/Files/AboutOT/consumers/Adults/LowVision/Low%20Vision%20Tip%20Sheet.pdf>

Technology Research to Assist Mobility for the Blind

72

- ❑ **Smart Phone App** - uses phone GPS, compass, accelerometers to provide highly accurate location information to locate crosswalks and building entrances
- ❑ **Radiofrequency Identification (RFID)** - technology may be incorporated into Braille signs to send information automatically to smartphones to read tags.
- ❑ **HandSight** will utilize a tiny camera embedded connected to a smartwatch to read text which is spoken out loud.

Copyright J. Gootkin 2019

Image <https://www.research.va.gov/currents/fall2014/fall2014-16.cfm>

Rehabilitation from Polytrauma Conditions

73

- Therapists may be involved in the management of musculoskeletal, neurologic, and mental health conditions as a result of combat or across the lifespan of the Veteran.
 - Amputation
 - Spinal Cord Injury
 - Traumatic Brain Injury
 - Post Traumatic Stress Disorder
 - Chronic Pain

Copyright J. Gootkin 2019

Understanding Combat Related Amputations

74

- Analysis of injury patterns and deaths during OEF and OIF indicates that most combat-related injuries occur as a result of injury from explosions, followed by gunshot wounds.
 - 78% Explosions
 - 18% Gunshot


Copyright J. Gootkin 2019

<https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFull.pdf>

Blast Casualty Injury Patterns

75

- Because of current advances in body armor, injury distribution is primarily to the extremities



Copyright J. Gootkin 2019

<https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFull.pdf>

Extremity Injury Resulting in Amputation

76

- Blast waves can shatter bones, avulse an extremity, cause extensive soft tissue damage, and flying debris create fragmentation or penetrating extremity injuries.

Copyright J. Gootkin 2019

Traumatic Amputation Immediate Medical Management

77

- Managing hemorrhaging with a proximal tourniquet while in the battlefield is critical to preserve life and avoid the need for massive transfusion.
- For high injuries with pelvic fracture, application of a Junctional Emergency Treatment Tool (JETT) is used to apply direct pressure.

Copyright J. Gootkin 2019

<https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFull.pdf>

Consider This

Battlefield Management of Amputations

78

- Procedure is to ligate vascular structures proximal to the bone resection, preserve bone length in absence of viable soft tissue, preserve atypical skin and tissue flaps, and leaving traumatic amputation wound open.
- Splints and bivalve casts stabilize the region with skin traction to avoid soft tissue retraction through the evacuation chain.

Copyright J. Gootkin 2019

Follow-up Medical Management of Amputation

79

- Negative Pressure Wound Therapy (NPWT) is used as a bridge to delayed closer, flap coverage, or coverage with a split thickness skin graft.
- It contributes to increased granulation rate, decreased bacterial colonization, and effective closure of the wound.

http://www.usaisr.amedd.army.mil/cpgs/Amputation_01Jul2016.pdf

Copyright J. Gootkin 2019

Residual Limb Wound Closure Techniques

80

- After staged wound debridement suturing or vessel loops can be stapled in a zig-zag pattern across the wound with some pressure applied.
- This aids closure by keeping the skin from passively retracting while also allowing granulation and mild drainage before grafting.

Copyright J. Gootkin 2019

Residual Limb Wound Closure Continued

81

- Scar patterns may be different and more extensive than traditional civilian amputations.

Copyright J. Gootkin 2019

Rehabilitation Outcomes

82

- Veterans with upper extremity amputations reported lower psychosocial adjustment, physical abilities, and prosthetic satisfaction than those with lower extremity amputations.

	Upper Extremity Amputation	Lower Extremity Amputation
Agree "life is full"	80%	91%
Agree "had gotten used to wearing an artificial limb"	23%	55%
Functional Outcome	Quick DASH 36.6/100	TUG 10.5 seconds

Copyright J. Gootkin 2019

<https://www.va.gov/oig/pubs/NAOIG-11-02138-116.pdf>

Upper Extremity Prosthetics Advances

83

- 2017 the VA unveiled LUKE – Life Under Kinetic Evolution -Arm that allows movement as one unit as opposed to traditional UE prosthetics where each joint is controlled individually through isolated movements.

Copyright J. Gootkin 2019

<https://www.aptcenr.research.va.gov/programs/ortho/natural-sensation-for-amputees/>

Lower Extremity Prosthetics Advances

84

- Osseointegration - A titanium coated implant is firmly anchored in living bone.
- The bone of the residual limb grows into the implant, and the prosthetic is attached directly to the metal connector of the implant without the need for a socket.

Copyright J. Gootkin 2019

Images https://www.amputation.research.va.gov/prosthetic_engineering/prosthetic_engineering_overview.asp

Lower Extremity Prosthetics Advances

85

- ❑ Powered Ankle Foot Prosthesis - ankle uses tendon like springs and an electric motor to replace lost muscle power and ankle/foot movement for propulsion during gait.
- ❑ Benefits include
 - ❑ Decreased energy expenditure
 - ❑ Improved balance
 - ❑ Decreased proximal joint damage
 - ❑ Improved gait pattern and walking speed

Au, S. K., Herr, H., Weber, J., & Martinez-Villalobos, E. C. (2007, August). Powered ankle-foot prosthesis for the improvement of amputee ambulation. In *Engineering in Medicine and Biology Society, 2007. EMBS 2007. 29th Annual International Conference of the IEEE* (pp. 3020-3026). IEEE.

Herr, H. M., & Grabowski, A. M. (2012, February). Bionic ankle-foot prosthesis normalizes walking gait for persons with leg Image Copyright © J. Gootkin 2019. https://www.amputation_research.va.gov/

Prosthetics Research

86

- ❑ Compliant Torque Adapter – located in transtibial pylon to allow easier maneuvering around obstacles.
- ❑ Stochastic Resonance – subthreshold vibration sensors in the socket may enhance peripheral sensation to improve postural stability and locomotion.
- ❑ Distributed Sensing – A flexible array built into the socket to measure moisture, temperature, pressure, and shear stress.

Images https://www.amputation_research.va.gov/prosthetic_engineering/prosthetic_engineering_overview.asp
Copyright J. Gootkin 2019

Prosthetic Research Continued

87

- ❑ Natural Sensation feedback is being investigated where sensors in the prosthetic hand control stimulation to nerves in the residual limb for tactile perception.

Copyright J. Gootkin 2019 <https://www.apccenter.research.va.gov/programs/ortho/natural-sensation-for-amputees/>
<https://www.research.va.gov/pubs/varqu/winter2016/11.cfm>

Smart Cane Research

88

- ❑ A cane equipped with a microprocessor and biofeedback control can measure cane contact with ground, gait cycle, and load phasing.
- ❑ Providing the user a sound or vibration signal from the cane will help them apply the optimal amount of force on the cane to reduce pain, decrease joint loading, and improve gait.

Copyright J. Gootkin 2019

Spinal Cord Injury (SCI)

89

- ❑ Gunfire, explosions, motor vehicle accidents and falls are the primary mechanisms of spinal cord injury in war zone.
- ❑ Spinal stabilization in the combat setting becomes challenging due to tactical considerations, body armor, and moving the injured thorough the evacuation chain.
- ❑ Main goal is spinal immobilization and spinal cord resuscitation to minimize secondary injury.
- ❑ Medial and surgical management mirrors civilian care

Copyright J. Gootkin 2019

SCI Mobility Research

90

- ❑ Advanced Lower Extremity Neuroprosthesis (LE-IST) - involves surgical implantation of neural stimulation technology with nerve or muscle electrodes activating muscles to allow standing.
- ❑ Similar technology is aimed at improving ambulation for individuals with incomplete spinal cord injury.

Copyright J. Gootkin 2019 https://www.apccenter.research.va.gov/studies/standing_transfer/index.asp
<https://www.apccenter.research.va.gov/studies/walking/index.asp>

SCI Mobility Research Continued

91

- Clinical trials are underway of an implanted trunk neuroprosthesis to control seated posture and balance for improved bed mobility and pulmonary function.

Copyright J. Gootkin 2019 [himages https://www.apccenter.research.va.gov/studies/trunk_control/index.asp](https://www.apccenter.research.va.gov/studies/trunk_control/index.asp)

SCI Mobility Research Continued

92

- **BrainGate** -uses implanted microelectrodes in the portions of the cortex that control voluntary movement.
 - Neural signals are transmitted to an external decoder that translates the into commands for electronic or robotic devices to restore movement.

Copyright J. Gootkin 2019 <https://www.research.va.gov/currents/1015-3.cfm>

SCI Mobility Research Continued

93

- Weight bearing is important for bone integrity, bowel function, and cardio-metabolic health.
- **ReWalk** - wearable robotic exoskeleton that provides powered hip and knee motion for standing, gait, and stair negotiation.

<https://www.research.va.gov/publications/varqu/winter2016/14.cfm>

Copyright J. Gootkin 2019

Traumatic Brain Injury

94

- "Signature Injury" of the current military conflict.

Copyright J. Gootkin 2019

dvbic.dcoe.mil/dod-worldwide-numbers-tbi

Blast Related TBI Mechanism of Injury

95

- Kinetic energy from blast explosions create massive wave that induce atmospheric pressure changes theorized to result in shearing of the axons, air embolism, hemorrhages, chromatolysis, and impaired vascular homeostasis.
- With closed head injury coup-counter coup injuries creates shearing forces and injury to the brain in regions deep and opposite to the site of impact.

Copyright J. Gootkin 2019 <https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFull.pdf>

Penetrating TBI Mechanism of Injury

96

- Damage can also occur from ballistic trauma where objects penetrate the skull.

Copyright J. Gootkin 2019

<https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFull.pdf>

TBI Medical Management 97

- ❑ The immediate goal is to minimize secondary brain injury by optimizing cerebral blood flow, mitigating elevations in intracranial pressure (ICP), and addressing intracranial hemorrhage and cerebral edema.
- ❑ The goal of the first 72 hours of care is to provide clinical stability, arrest ongoing injury, preserve neurological function, and prevent secondary medical complications.
- ❑ Poor outcomes associate with systolic BP below 90 mm Hg and ICP over 25 mm Hg.

Copyright J. Gootkin 2019 <https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFULL.pdf>

Department of Defense TBI Case Definitions 98

- Penetrating TBI**
 - Scalp, skull and dura mater are penetrated.
- Severe TBI**
 - Confused/disoriented more than 24 hours or LOC more than 24 hours or memory loss more than 7 days.
- Moderate TBI**
 - Confused/disoriented more than 24 hours or LOC more than 30 minutes but less than 24 hours or memory loss greater than 24 hours but less than 7 days.
- Concussion/Mild TBI**
 - Confused/disoriented less than 24 hours or loss of consciousness up to 30 minutes or memory loss lasting less than 24 hours.

Copyright J. Gootkin 2019 Consider This

TBI Clinical Presentation Glasgow Coma Scale (GCS) 99

❑ Assesses level of consciousness through eye opening, verbal response, and motor.

BEST EYE OPENING (E)		BEST VERBAL RESPONSE (V)		BEST MOTOR RESPONSE (M)	
		Oriented, alert	5	Follows commands	6
Eyes open spontaneously	4	Confused, appropriate	4	Localizes to pain	5
Eyes open to speech	3	Disoriented, inappropriate	3	Withdrawal to pain	4
Eyes open to pain	2	Incomprehensible speech	2	Flexor posturing	3
No response	1	No response	1	Extensor posturing	2
		No response	1	No response	1

Mild TBI
13 or greater

Moderate TBI
9 to 12

Severe TBI
8 or less

Copyright J. Gootkin 2019 <https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFULL.pdf>

Mild TBI (mTBI) Clinical Presentation 100

- Physical**
 - Headache, fatigue, dizziness, lethargy
- Cognitive**
 - Easily confused/distracted, difficulty with memory, attention, concentration, and problem solving.
- Sensory**
 - Impaired vision, hearing, taste, smell, or touch
- Behavioral**
 - Depression anxiety, agitation, irritability, anger, sleep disturbance

Copyright J. Gootkin 2019

mTBI Return to Activity Progression 101

Stage 1 Rest

Stage 2

Light Routine Activity
30 mins physical activity then 4 hours rest
Max 30 mins cognitive activity then 60 mins rest

Stage 3

Light Occupation Activity
60 mins physical activity then 4 hours rest
Max 30 mins cognitive activity then 60 mins rest

Stage 4

Moderate Activity
Max 90 mins activity at 12-16 RPE
20 mins cognitive activity then 40 mins rest

Stage 5

Intensive Activity
Resume usual exercise PRE 16+
50 mins cognitive activity at a time

Stage 6 Unrestricted Activity

Progress every 24 hrs if asymptomatic

Copyright J. Gootkin 2019
Department of Defense. (2014). Progressive return to activity following acute concussion/mild TBI: guidance for the rehabilitation provider in deployed and non-deployed settings. Clinical Recommendation. Clinical Recommendation January 2014.

mTBI Post-Concussive Syndrome 102

- ❑ Residual symptoms of headache, dizziness, irritability, decreased concentration, photosensitivity, fatigue, sleep disturbance, attention/concentration impairments and forgetfulness following mild traumatic brain injury for greater than three months.
- ❑ Management is typically symptom based with majority of individuals reporting headache as the primary persistent symptom.

Copyright J. Gootkin 2019

TBI Rehabilitation

103

- For mTBI, treatment is symptom specific and multi-modal monitoring for various system involvement and directing referral to the appropriate medical professional for additional assessment.
 - Concussion monitoring
 - Post-traumatic headaches
 - Sensory Impairments
 - Neuroendocrine dysfunction

Copyright J. Gootkin 2019

Rehabilitation Concussion Self Monitoring

104

- Developed by the VA National Center and National Center for Telehealth & Technology, the Concussion Coach app allows tracking of TBI symptoms and resources.
 - Self-assessment of emotional changes, cognitive issues and physical problems
 - Relaxation exercises to manage symptoms
 - Immediate access to crisis resources, personal support and professional mental health care.

Copyright J. Gootkin 2019 <http://t2health.dcoe.mil/apps/ConcussionCoach>

Rehabilitation Post-traumatic Headaches

105

	Tension	Cervicogenic	Migraine
Pain	Mild -moderate Tighten, pressure	Mild -moderate Tighten. burn	Severe- debilitating Throbbing, pulsing
Location	Bilateral frontal and posterior cervical	Unilateral cervical and occipital	Unilateral frontotemporal or retro-orbital
Duration	30 mins - 7 days	Variable	4-72 hours
Phono/ photophobia	No	No	One or both
Function	Normal	Normal	Impaired
Nausea	No	No	Yes
Tenderness	Facial and neck	Posterior neck	Not typical
Medication	Abortive NSAID OTC analgesic	Abortive NSAID	Abortive or Prophylactic

Copyright J. Gootkin 2019

Rehabilitation Sensory Impairments

106

- Visual acuity is typically intact, but ocular misalignment, and poor eye coordination are associated with patient reports of difficulty focusing, blurred or double vision impairing reading and gait.
- Despite normal audiology testing, Veterans report problems hearing speech in difficult listening environments indicating damage to the central auditory system.

Copyright J. Gootkin 2019 <https://www.research.va.gov/topics/tbi.cfm>

Rehabilitation Neuroendocrine Dysfunction

107

- With brain injury vascular damage, pituitary inflammation, or a biochemical response to blast exposure may interfere with normal hormone production and regulation by the pituitary and hypothalamus.
- Example: In female symptoms may include excessive fatigue, low muscle mass, increased body fat around the waist, low blood pressure, and amenorrhea.
- Referral to endocrinologist for hormonal screening is warranted.

Copyright J. Gootkin 2019 http://dvbic.dcoe.mil/files/resources/DCoE_TBI_NED_Training_Slides.pdf

Consider This

Traumatic Brain Injury Research

108

- **Cerebellar Trauma** – Repeated blast exposures ruptures part of the blood-brain barrier leading to loss of neurons and damage to the cerebellum.
- **Chronic Traumatic Encephalopathy (CTE)** – Similar to athletes, post-mortem analysis of Veterans who sustained repetitive concussions revealed degenerative CTE.

Copyright J. Gootkin 2019 <https://www.research.va.gov/topics/tbi.cfm>

TBI Research Continued

109

- **Alzheimer's Disease** – Veterans 55 and older with a past diagnosis of TBI had a 60% greater risk of developing dementia.
- **Brain Aging** – Exposure to blasts without resulting concussion may still lead to fraying of brain white matter with premature brain aging.

<https://www.research.va.gov/topics/tbi.cfm>

Copyright J. Gootkin 2019

Post-Traumatic Stress Disorder (PTSD)

110

- PTSD is a psychological condition that affects those who have experienced a traumatizing or life-threatening event such as combat, natural disasters, serious accidents, or violent personal assaults.

<https://www.ptsd.va.gov/>
<https://ia902709.us.archive.org/25/items/CombatCasualtyCare/CCCFull.pdf>

Copyright J. Gootkin 2019

Psychological First Aid (PFA)

111

- Appropriately supporting individuals in the immediate aftermath traumatic experiences helps reduce the initial stress they are experiencing and assists in developing long and short term adaptive functioning skills.

Response	What to day and do
Concern, shame, guilt, fear	Reminders that you are not "going crazy". If feelings persist for a month seek care.
Overwhelmed by tasks	Identify priorities, make lists, rely on others
Fears of recurrence and reactions to reminders	Triggers can be sounds, smells, time of day, media coverage.

https://www.ptsd.va.gov/professional/materials/esset/pfa_mobile_app.asp

Copyright J. Gootkin 2019

Disaster and Terrorism

112

- The goal is to protect survivors from further harm, reduce psychological arousal, provide information and foster communication, keep families together and facilitate reunion with loved ones.

https://www.ptsd.va.gov/professional/materials/esset/pfa_mobile_app.asp

Copyright J. Gootkin 2019

PTSD Progression

113

The diagram illustrates the progression of PTSD over time. It starts with a starburst labeled 'Trauma'. A horizontal arrow points to the right, with three time markers: '2 Days', '1 Month', and '3 Months'. Below the arrow, four stages are listed: 'Acute/Combat Stress (ASR/COSR)' under 2 Days, 'Acute Stress Disorder (ASD)' under 1 Month, 'Acute PTSD' under 3 Months, and 'Chronic PTSD' under 3 Months.

If the body's stress and coping systems do not manage the trauma, a complex cluster of symptoms results.

Department of Defense (2010) Clinical Practice Guideline for the Management of Post Traumatic Stress Disorder. *Guideline Summary 2010*.

Copyright J. Gootkin 2019

PTSD Symptoms

114

- Last more than one month after trauma and lead to significant impairment in ability to interact in social, occupational, or other important areas of function.
- Individual demonstrates a combination of physical, mental, and behavioral symptoms in addition to changes in functioning.

Copyright J. Gootkin 2019

PTSD Common Presenting Symptoms 115

- Physical**
 - Chronic Pain, Migraines, Vague Somatic Complaints
- Mental**
 - Poor Attention and Abstract Thinking, Anxiety, Depression
- Behavioral**
 - Irritability, Guilt, Avoidance, Anger, Noncompliance, Self-risk, Aggression, Threatening Behavior

Copyright J. Gootkin 2019

PTSD Symptom Clusters 116

- Re-Experiencing**
 - Intrusive memories, images, perceptions
 - Exaggerated emotional and physical reactions
- Avoidance & Emotional Numbing**
 - Activity avoidance
 - Detached
 - Restricted Emotion
- Increased Arousal**
 - Sleep disturbance
 - Irritability
 - Hypervigilance
 - Difficulty Concentrating

Copyright J. Gootkin 2019

PTSD Screening Tools 117

- ❑ Referral to a mental health provider may be necessary if symptoms persist and interfere with function and relationships.
- ❑ Only specifically trained mental health professionals can use advanced assessment tools.
 - ❑ Evaluation of Lifetime Stressors (ELS)
 - ❑ Life Events Checklist for DSM-5 (LEC-5)
 - ❑ Combat Exposure Scale (CES)

Copyright J. Gootkin 2019

PTSD Screening Tool Evaluation of Lifetime Stressors (ELS) 118

- ❑ Self-report questionnaire and interview assessing perceptions and emotional response to disaster, illness, accident, violence, combat and other trauma.
- ❑ The interview provides a way to explore dimensions of each trauma including trauma type, perpetrators/victims, duration, frequency, perceptions of threats and emotional response, and others.

Copyright J. Gootkin 2019

PTSD Screening Tool Life Events Checklist for DSM-5 (LEC-5) 119

- ❑ Self-report measure designed to screen for potentially traumatic events in a respondent's lifetime.
- ❑ Specifically assesses exposure to 16 events known to potentially result in PTSD or distress.

Copyright J. Gootkin 2019

PTSD Screening Tool Combat Exposure Scale (CES) 120

- ❑ 7-item self-report measure that assesses wartime stressors and exposures experienced by combatants. Items are rated on a multiple scales.
- ❑ Frequency
 - 1 = "no" or "never" to 5 = "more than 50 times"
- ❑ Duration
 - 1 = "never" to 5 = "more than 6 months"
- ❑ Degree of loss
 - 1 = "no one" to 4 = "more than 50%"

Copyright J. Gootkin 2019

PTSD Management 121

- Ensure basic physical needs are met
- Interventions to preserve life if individual is unstable or dangerous to self or others.
- Medical and Psychological Care
- Cognitive Behavioral Therapies
- Social and Spiritual Support
- Medications
- Cognitive Fitness and Psychological Resilience Strategies
- Relaxation and Stress Management

Copyright J. Gootkin 2019

Rehabilitation Considerations for Patients with PTSD 122

- Patients diagnosed with PTSD may be participating in therapy for rehabilitation from musculoskeletal, general medical, and neurologic conditions or to promote health and wellness to manage their PTSD symptoms.
- Therapists should perform basic assessment to understand the patients status and recognize how to manage therapeutic interactions.

Copyright J. Gootkin 2019

Screening Tool Primary Care PTSD (PC-PTSD) 123

Designed for healthcare professionals outside the mental health fields.

- **Would it be ok to talk about stress?**
In your life, have you ever had an experience so horrible, frightening, or upsetting that, in the past month you...
- Have had nightmares about it or thought about it when you did not want to?
- Tried hard not to think about it or went out of your way to avoid situations that reminded you of it?
- Were constantly on guard, watchful, or easily startled?
- Felt numb or detached from others, activities, or your surroundings?

Veterans Crisis Line 1-800-273-8255 (Press 1)
Copyright J. Gootkin 2019 <https://www.va.gov/oas/pocketcard/Military-Health-History-Card-for-print.pdf>

PTSD Managing Interactions 124

- Do not approach the individual from behind or sneak up on them
- When interviewing the patient focus on establishing a strong therapeutic alliance.
 - Ensure privacy.
 - Make no assumptions about feelings or meaning.
 - Show interest and concern.
- If patient is resistant to referrals, emphasize value of evaluation and treatment and suggest resources they can explore first on their own.

Copyright J. Gootkin 2019

PTSD Managing Interactions Continued 125

- Acknowledge distress offering empathy and validating patient's feelings.
- Example statements include:
 - "I am so sorry that you are struggling with this."
 - "I can appreciate how difficult this is for you."
 - "You are not the only patient I have who is struggling with this."
 - "It's not easy, is it?"

Copyright J. Gootkin 2019

PTSD Managing Interactions Continued 126

- If the patient experiences PTSD symptoms during therapy, the clinician should speak in a calm, matter of fact voice.
- Reassure the patient that everything is okay.
- Remind the patient that they are in a safe place and their care and well being are a top priority.
- Explain medical procedures and check with the patient (e.g., "Are you ok?").
- Ask (or remind) the patient where he or she is right now.
- If the patient is experiencing flashbacks, remind him or her that they are in XYZ office at a specific time in a specific place (grounding).
- Offer the patient something that will make them feel more like their usual self.

Copyright J. Gootkin 2019 https://www.ptsd.va.gov/professional/manuals/manual.pdf/lwca/lrca_clinician_guide_ch_7.pdf

Pain Syndromes

127

Pain is associated with trauma from combat injuries to multiple body systems combined with potential amplification of comorbidities of PTSD and TBI.

PTSD

Post-Deployment Multi-system Disorder (PMD)

TBI

Chronic Pain

Consider This

Copyright J. Gootkin 2019

Pain Patterns

128

- Some sources view Pain Level as the Fifth Vital Sign
- The most commonly reported pain is due to arthritis, back pain, neuropathic pain, and headaches.

Temperature

Heart Rate

Blood Pressure

Respiratory Rate

Pain Level

Copyright J. Gootkin 2019

Chronic Pain

129

- Following combat-related injuries, increased incidence of chronic pain greater than three months is associated with:
 - Ongoing sleep disturbance
 - Emotional distress
 - Residual traumatic brain injury symptoms
- Low pain self-efficacy, maladaptive coping strategies, reluctance to access mental health care may contribute to the cycle of pain.

Enhancing Veteran-centered care: a guide for nurses in non-VA settings: Johnson BS, Boudiab LD, Freundt W, Anthony W, Gienzek GB, Carter J. Am J Nurs. 2013 Jul; 113(7):24-39; quiz 54, 40.

Consider This

Copyright J. Gootkin 2019

Pain Screening Tools

130

- Standard musculoskeletal assessments should determine if a physiologic etiology for the pain exists.
- When diffuse somatic, cognitive, and emotional complaints are associated with pain multiple system polytrauma assessments should be utilized.

Copyright J. Gootkin 2019

Pain Screening Tool Defense and Veterans Pain Rating Scale (DVPRS)

131

Visual analog scale (VAS) with functional clarifiers and supplemental pain questions to evaluate biopsychosocial impact of pain.

https://www.va.gov/PAINMANAGEMENT/docs/DVPRS_2slides_and_references.pdf

Copyright J. Gootkin 2019

Pain Screening Tool DVPRS Clarifying Questions

132

- Circle one number that describes how, during the past 24 hours pain has:
 - interfered with you usual ACTIVITY
 - interfered with you SLEEP
 - affected your MOOD
 - contributed to your STRESS

Does not Completely

Copyright J. Gootkin 2019

Pain Pharmaceutical Management 133

- ❑ Condition specific non-opioid agents should be optimized before opiates are considered for short durations.
 - ❑ Risk of dependence, addiction, and death.
- ❑ Narcotics and muscle relaxants can slow cognition and executive functions.
- ❑ Complexity in symptom relief from narcotics needs to be recognized as reduction of psychological anxiety may be confused with diminished pain intensity.

Howell, P., Capehart, B. P., & Hoenig, H. (2015). Rehabilitation of combat-related injuries in the Veterans Administration A web of support. *North Carolina Medical Journal*, 76(5), 323-327. <https://www.healthquality.va.gov/guidelines/Pain/cot/VADoDOTCPGProviderSummary022817.pdf>

Copyright J. Gootkin 2019

Adverse Effects of Opioids 134

Respiratory Depression	• Drowsiness, slow shallow breathing, loud or unusual snoring
Mental Status Changes	• Confusion, hallucinations, restlessness, dysphoria, seizures, agitation
Opioid Induced Endocrinopathy	• Loss of libido, fatigue, mood alterations, loss of muscle mass/strength, abnormal menses
Sleep Apnea	• Loud snoring, morning headaches, irregular pauses in breathing depression/emotional instability

Copyright J. Gootkin 2019 <https://www.healthquality.va.gov/guidelines/Pain/cot/ManagingSideEffectsFactSheetFINAL2017.pdf>

Pain Non-Pharmaceutical Management 135

- ❑ Comprehensive management must address biological, psychological, and psychosocial aspects of pain.
- ❑ Distinguish pain from musculoskeletal dysfunction or neurologic alterations in tone.
- ❑ Biofeedback, exercise, thermal modalities
- ❑ Relaxation exercises and meditation
- ❑ Good sleep hygiene
- ❑ Patient education

Copyright J. Gootkin 2019

Pain Cognitive Behavioral Therapy 136

- ❑ Encouraging increasing activity by setting goals
- ❑ Correcting false and unrealistic beliefs/concerns about pain
- ❑ Teaching cognitive and behavioral coping skills (e.g., activity pacing)
- ❑ Practicing and consolidation of coping skills and reinforcement of use.

Copyright J. Gootkin 2019

Non-Somatic Disorders 137

- ❑ Post deployment individuals experience a variety of physical conditions (somatic) in addition to presenting with less tangible non-somatic mental health conditions that contribute to the complex management of polytrauma.
 - ❑ Anxiety
 - ❑ Drug Use
 - ❑ Sleep Disturbance

Copyright J. Gootkin 2019

Anxiety Assessment Beck Anxiety Inventory (BAI) 138

- ❑ The 21-item self-report measures emotional, physiological, and cognitive symptoms of anxiety.
- ❑ Respondents rank how bothered they are by each of the symptoms.

0-7 Minimal	8-15 Mild	16-25 Moderate	26-63 Severe
-------------	-----------	----------------	--------------

Copyright J. Gootkin 2019

Anxiety Assessment in Polytrauma BAI Continued

139

- Symptoms can be aligned to specific factors associated with anxiety measured with the BAI.
- Safety threat component provides information regarding the mental health of combat Veterans.

Subjective Fear
• Heart pounding

Neurophysiological
• Unsteady

Panic
• Shaky

Somatic
• Face flushed

Safety threat
• Fear of dying

Consider this

Palmer, G. A., Happe, M. C., Paxson, J. M., Jurek, B. K., & Olson, S. A. (2016). Dimensionality of the Beck Anxiety Inventory in a polytrauma sample of OEF/OIF/OND Veterans: Findings from exploratory and confirmatory factor analysis. *Military Psychology, 28*(3), 206.

Copyright J. Gootkin 2019

Stress, Trauma, Alcohol, and Drug Use Assessment

140

- Drug and alcohol use is common to self-medicate.
- Common statements attempting to self-assure there is not a problem with dependency or overuse of drugs or alcohol.
 - "I just drink beer (wine)"
 - "I don't use hard drugs"
 - "I'm not an alcoholic"
 - "I gave it up for 3 weeks last year"
 - "I don't drink every day"
 - "I've never missed a day of work"
 - "I don't need help, I can handle it myself"

https://www.ptsd.va.gov/professional/manuals/manual-pdf/iwecg/iraq_clinician_guid_e_app_14.pdf

Copyright J. Gootkin 2019

Traumatic Stress in Female Veterans

141

- As of 2015, women are being integrated into military combat positions and receiving the specialized training that they did not receive in prior years.
- Changing theaters of operation may contribute to feeling along compounded by worrying about family at home during long deployments.
- High social and emotional support after war has been shown to contribute to lower stress levels.

<https://www.defense.gov/News/Article/Article/648766/officials-describe-plans-to-integrate-women-into-combat-roles/>
https://www.ptsd.va.gov/professional/trauma/war/traumatic_stress_in_female_veterans.asp

Copyright J. Gootkin 2019

Sleep Disturbance

142

- Difficulty initiating or maintaining sleep and non-restful sleep may be related to experiences during deployment for combat or peacekeeping missions.
 - Nightmares result in waking up feeling terrified.
 - Saying awake to "stand guard" rather than sleep.
 - Irregular sleep-wake cycles from shifts and loud noise exposure.

Copyright J. Gootkin 2019

Sleep Disturbance and Comorbidities

143

Theories:

- Altered hypothalamus neuropeptide
- Altered melatonin production by pineal gland
- Intrusive chronic headaches

mTBI Post Concussion

Sleep Disturbance

Pain

Impaired Cognition

Irritability

Association with PTSD

Quellet, M. C., Beaulieu-Bonneau, S., & Morin, C. M. (2015). Sleep-wake disturbances after traumatic brain injury. *The Lancet Neurology, 14*(7), 746-757.

Copyright J. Gootkin 2019

Sleep Disturbance Assessment

144

- Defense Centers of Excellence recommends including an interview question regarding sleep:
 - Are you experiencing frequent difficulty falling or staying asleep, excessive daytime sleepiness or unusual events during sleep?
- Pittsburgh Sleep Quality Index (PSQI) is a self-report with higher global score on the 0-21 scale indicating worse sleep quality.
 - Validated on TBI patients with >5 indicating sleep disturbance

Department of Defense. (2014). Management of sleep disturbance following concussion/mild TBI: guidance for primary care management in deployed and non-deployed settings. *Clinical Recommendation*. June 2014.

Copyright J. Gootkin 2019 <http://www.psychiatry.pitt.edu/research/tools-research/assessment-instruments/pittsburgh-sleep-quality/>

Sleep Hygiene and Stimulus Control

145

- Keep bed only for sleep
- Decrease stimulants and alcohol
- Have a regular bedtime and rising time
- Get exposure to sunlight every morning
- Increase regular exercise
- If you don't fall asleep within 30 minutes, get up
- Wind down before bedtime
- Remove electronics your bedroom
- Go to bed only when sleepy

Copyright J. Gootkin 2019

Compassion Fatigue

146

- Providing care and support to service members and Veterans can lead to burnout and secondary traumatic stress for the healthcare provider.
- The Provider Resilience app created by National Center for Telehealth & Technology includes tools for monitoring to avoid compassion fatigue.
 - Self assessments
 - Inspirational videos
 - Tools for restful breaks

Copyright J. Gootkin 2019 <http://t2health.dcoe.mil/apps/provider-resilience>

Conclusion

147

- Remember all that our men and women in uniform have selflessly sacrificed.
- Analyze their conditions from every perspective and be an advocate to help them access the extensive VA resources available to foster the independence and recovery they deserve.

Copyright J. Gootkin 2019 <https://www.polytrauma.va.gov/>

Combat-Related Polytrauma Resource Links

Veterans Administration Medical Services

Veterans Administration Medical and Benefits Center Locations

https://www.va.gov/landing2_locations.htm

State Veterans Affairs Offices <https://www.va.gov/statedva.htm>

Veterans Choice Program: Information to become a provider

https://www.va.gov/opa/choiceact/documents/How_to_Become_VA_Provider_05242017_508.pdf

VA Polytrauma/TBI System of Care: Resources and Services <https://www.polytrauma.va.gov/>

VA Amputation System of Care <https://www.prosthetics.va.gov/asoc/index.asp>

VA Spinal Cord Injuries and Disorders System of Care <https://www.sci.va.gov/>

VA Substance Abuse Programs <https://www.mentalhealth.va.gov/res-vatreatmentprograms.asp>VA

Embedded Fragments Surveillance Center: Urine Biomonitoring for potential toxicity

https://www.publichealth.va.gov/exposures/toxic_fragments/surv_center.asp

VA Blind Rehabilitation Services: Low vision care, inpatient rehabilitation, and blind services

https://www.prosthetics.va.gov/blindrehab/BRS_Coordinated_Care.asp

Veterans Administration Social Services

Homeless Veterans Resources and Referral Information <https://www.va.gov/HOMELESS/index.asp>

Homeless Veterans Outreach & Education Tools https://www.va.gov/HOMELESS/get_involved.asp

Project Hired Wounded Warrior Workforce: Career Exploration, Job Placement, and Transition Support <http://www.projecthired.org/jobseekers/veterans>

Veterans Service Organizations: Resources for Personnel, Veterans, and Families

<https://www.ptsd.va.gov/professional/materials/web-resources/military-resources.asp>

Veterans Administration Mental Health

Veterans Crisis Line: 1-800-273-8655 Press 1 <https://www.veteranscrisisline.net/>

VA Mental Health Self-Help Resources: https://www.mentalhealth.va.gov/self_help.asp

VA National Center for PTSD: <https://www.ptsd.va.gov/>

VA Mental Health Community Provider Toolkit: Multiple resources and free consultation service for clinicians managing patients with PTSD

<https://www.mentalhealth.va.gov/communityproviders/index.asp>

Smartphone Apps

VA Psychological First Aid Mobile: Resources and tips on applying Psychological First Aid
<http://t2health.dcoe.mil/apps/PFAMobile>

National Center for Telehealth & Technology T2 Provider Resilience: Tools to monitor and guard against burnout and compassion fatigue <http://t2health.dcoe.mil/apps/provider-resilience>

National Center for Telehealth & Technology T2 Mood Tracker: Track Emotional Health
<http://t2health.dcoe.mil/apps/t2-mood-tracker>

National Center for Telehealth & Technology T2 Concussion Tracker: Self Assessment of Symptoms and Access to Crisis Resources <http://t2health.dcoe.mil/apps/ConcussionCoach>

Screening Tools

VA Military Trauma Iran War Clinical Guide
<https://www.ptsd.va.gov/professional/materials/manuals/iraq-war-clinician-guide.asp>

Department of Defense Dictionary of Military and Associated Terms:
http://www.dtic.mil/doctrine/dod_dictionary

VA Military Culture Training for Health Care Professionals <https://health.mil/Training-Center/DoD/Military-Culture-Training-for-Health-Care-Professionals--Introduction-and-Self-Awareness-Exercise>

VA Military Health History Card: Patient Interview Questions
<https://www.va.gov/oaa/pocketcard/Military-Health-History-Card-for-print.pdf>

VA Psychological First Aid Manual Field Guide
<https://www.ptsd.va.gov/professional/materials/manuals/psych-first-aid.asp>

Beck Anxiety Inventory <http://www.pearsonclinical.com/psychology/products/100000251/beck-anxiety-inventory-bai.html?Pid=015-8018-400&Mode=summary>

Vision Screening: Convergence Insufficiency Symptom Survey
<http://www.sankaranethralaya.org/pdf/patient-care/Convergence-Insufficiency-Symptom-Survey.pdf>

Defense and Veterans Pain Rating Scale
https://www.va.gov/PAINMANAGEMENT/docs/DVPRS_2slides_and_references.pdf

Primary Care PTSD Screen
https://www.mentalhealth.va.gov/communityproviders/docs/PCPTSD_Screen.pdf

Pittsburgh Sleep Quality Index <http://www.psychiatry.pitt.edu/research/tools-research/assessment-instruments/pittsburgh-sleep-quality-index-psqi>